## Electronic Structure Studies with Spin-Orbit Coupling Effect of the Molecule TII

GHINA CHAMIEH, LOKMAN AWAD, MAHMOUD KOREK<sup>\*</sup> Faculty of Science, Beirut Arab University, P. O. Box 11-5020 Riad El Solh, Beirut 1107 2809, LEBANON

#### \*Corresponding Author

Abstract: - To study the low–lying electronic states of the TII molecule, the electronic structure of this molecule has been investigated via an ab initio Complete Active Space Self Consistent Field and the Multireference Configuration Interaction with Davidson correction calculation (CASSCF/MRCI+Q). In the representations of  $^{2s+1}\Lambda^{(+/-)}$  and  $\Omega^{(\pm)}$ , the adiabatic potential energy curves (PECs) along with static and transition dipole moment (DM) curves for 19 low-lying electronic states for TII molecule have been investigated. For the low-lying electronic states of this molecule, the spectroscopic constants Re,  $T_e$ ,  $\omega e$ , and Be, are provided. Based on the data obtained, this molecule is not a candidate for a Doppler laser cooling study.

*Key-Words:* - ab initio calculation, electronic structure, spin-orbit coupling, potential energy curves, dipole moment, spectroscopic constants, diatomic molecule.

Received: March 15, 2023. Revised: October 17, 2023. Accepted: November 29, 2023. Published: December 31, 2023.

## **1** Introduction

With a spin-orbit splitting of 7793 cm<sup>-1</sup>, thallium exhibits strong relativistic effects as a heavy metal belonging to the GIII-A group, which are anticipated to be significant for chemical bonding, [1]. It is well known that thallium compounds make valuable materials, such as high-temperature superconducting materials and catalysts for organic synthesis, [2], [3]. Thallium halides have drawn much attention from researchers because they are among the earliest thallium compounds to be produced. The spectra and dynamics of the electronic states of these compounds have been the subject of several experimental and theoretical studies.

Here, we use gaseous thallium mono-halide as representations straightforward of thallium compounds. Both theorists and experimentalists have long been drawn to these systems. For TIX (X = F, Cl, Br, and I), several spectroscopic measurements have been made up to this point, [4]. However, most excited states' geometrical and spectroscopic characteristics are yet unknown. The ground state and some excited electronic states of TlI were investigated, [4]. Spin-orbit coupling (SOC) significantly impacts spectroscopy and dynamics. The SOC effect can cause predissociations between  $\Lambda$ -S states with various spin

multiplicities and multiple avoided crossings of  $\Omega$  <sup>(±)</sup> states, [5].

The present work on the molecule TII is a part of our research project on iodine compounds such as CaI, [6], BaI, [7], MgI, [8] and BeI, [9]. Since there is some available data for electronic states in the literature on the molecule TII, we expect to supply pleasant details for theoretical studies of this molecule, [4]. Our powerful motive is to profoundly analyze the molecule's ground with excited states with their complete spectroscopic and spin-orbit studies in the current work.

Because of this absence, an ab initio method with a Complete Active Space Self Consistent Field has been used with Davidson correction (CASSCF/ MRCI+Q) to calculate the potential energy (PECs), the static and transition dipole moment curves (DMCs), the Franck-Condon factor FCF, and the radiative lifetime. In the representations,  $^{2s+1}\Lambda^{(+/-)}$ and  $\Omega^{(\pm)}$ , 19 low-lying electronic states have been investigated for the molecule of TII where most of the investigated data are presented here for the first time.

## 2 Computational Methods

The state average Complete Active Self Consistent Field (CASSCF)/Multireference Configuration Interaction (MRCI + Q) has been used to investigate the singlet and triplet electronic states of the TII molecule. With the graphical interface GABEDIT, [10]. The computational chemistry program MOLPRO (a software package developed for accurate ab initio quantum chemistry calculations) is used to accomplish these calculations, [11].

The Born-Oppenheimer approximation was used to obtain the potential energy curves. Using the customized core polarization potentials (CPP) and energy-consistent effective core potentials (ECP) from the Stuttgart/Cologne Group, the relativistic effects and core electrons were treated. The MRCI and CASSCF methods, along with selected basis sets, were used to describe the valence electrons.

For the TII molecule, the ECP60MDF basis set is used for the Tl atom with 21 valence electrons distributed as  $5s^2 5p^6 5d^{10} 6s^2 6p^1$ , [12]. For the I atom, the basis set ECP46MDF is considered with seven valence electrons distributed as  $5s^2 5p^5$ , [13]. For the considered molecule TII, 9 occupied shells are investigated in symmetry 1, 4 in symmetries (2 and 3), and only 1 shell in symmetry 4. For the closed shells, five are considered in symmetry 1, 2 in symmetries (2 and 3), and 1 shell in symmetry 4. We chose ECP as a basis for Tl and I atoms by referring to many papers on these atoms published in the literature, [14], [15], [16].

The diatomic molecules belong to  $C_{\infty v}$ , and because of the limitation of the MOLPRO, it turns to its subgroup  $C_{2y}$ , which has four irreducible representations (a1, b1, b2, a2). The active space for the TII molecule is  $4\sigma$  (TI: 6p0, 7s; I: 5p0, 6s),  $2\pi$ (Tl:  $6p\pm 1$ ; I:  $5p\pm 1$ ;) and  $0\delta$ , with the irreducible representation 7a1, 3b1, 3b2, and 1a2 noted by [4, 2, 2, 0]. To obtain the potential energy curves, the estimated energy points are connected using the avoided-crossing rule for electronic states that belong to the same irreducible representation of the single/double point group  $C\infty v$ . The onedimensional Born-Oppenheimer Schrödinger equation is used to obtain the spectroscopic constants including R<sub>e</sub> (equilibrium bond length), T<sub>e</sub> (transition energy),  $\omega_e$  (harmonic constant), and  $B_e$ (rotational constant).

### **3** Results and Discussion

#### **3.1 Spectroscopic Constants**

With and without spin-orbit coupling, the singlet and triplet electronic states of the molecule TII are investigated. The spectroscopic constants (Transition energies, internuclear distances, harmonic vibrational frequencies) for the bound states can be computed using the X-Poly or X-min programs. The PECs are first fitted to a polynomial function (U(r) =  $a_0 + a_1r + a_2r^2 +...$ ) to reproduce the ab initio values. After that, the constants are computed analytically by their definitions.

The operating system that is being used determines which program is selected. The Windows operating system is used with X-poly, and the Ubuntu operating system is used with X-min. However, both provide the same reliability and accuracy. Table 1 and Table 2 present the PECs spectroscopic constants that we computed using the X-poly program for the different electronic states.

Table 1. Spectroscopic constants  $T_e$ ,  $R_e$ ,  $\omega_e$ , and  $B_e$  of the molecule TII (with spin-orbit coupling)

states	T <sub>e</sub> (cm <sup>-1</sup> )	$\omega_e ({ m cm}^{-1})$	$R_{e}(\dot{A})$	B <sub>e</sub> (cm <sup>-1</sup> )
$(1)^{1}\Sigma^{+}_{0+}$	0.00	141.64	5.4712	0.0257
$(1)^{3}\Pi_{0+}$	19506.20	20.37	4.2567	0.0119
$(1)^{3}\Delta_{1}$	24169.04	29.70	4.0363	0.0132
$(1)^{3}\Sigma^{+}_{0+}$	24463.03	20.86	4.2342	0.0120
$(1)^3\Pi_1$	24696.92	19.78	5.4815	0.0071
$(1)^{3}\Sigma^{+}_{1}$	24758.26	15.71	5.9989	0.0601
$(1)^{3}\Delta_{2}$	24942.12	49.20	4.8681	0.0091
$(2)^{3}\Delta_{1}$	29702.54	23.34	4.2024	0.0122

Table 2. Spectroscopic constants  $T_e$ ,  $R_e$ ,  $\omega_e$  and  $B_e$  of the molecule TII (without spin-orbit coupling)

State	T.(cm-3)	04 (cm-1)	Re (Å)	B. (cm <sup>-1</sup> )
$\mathbf{X}^{1}\Sigma^{*}$	0.00 * 0.00 *	140.22* 151.00 *	2.9209* 2.823b	0.0252+
<b>(1)</b> <sup>3</sup> Σ*	23391.31 * 27493 *	40.15* 46.00*	4.100* 3.599*	0.0125
(1) <sup>9</sup> Δ	23730.11* 27971.00*	38.90* 41.00*	3.7360* 3.694*	0.01534
$(3)^{1}\Delta$	23734.06	22.44	4.3691	0.0113
<b>(1)</b> <sup>2</sup> Δ	23763.11 = 28042.00 =	37.55 4 40.00 *	4.2173 * 3.748 *	0.1103
(2) <sup>1</sup> Δ	23763.29=	21.99*	4.4192 =	0.01094
$(1)^{2}\Sigma^{-}$	23881.79=	11.86*	4.6612 =	0.0098
(2) <sup>3</sup> Σ°	23886.75*	10.24 *	4.9365*	0.0088
(1) <sup>3</sup> Σ <sup>-</sup>	23888.61 *	12.63+	4.6182=	0.0101
(2) <sup>5</sup> Δ	23945.91=	29.99*	7.0189*	0.0044
(3) <sup>2</sup> Σ*	25724.64*	38.89*	6.1775=	0.0056
(3) <sup>3</sup> ∏	42713.81=	38.91 *	3.7368=	0.0154
(3) <sup>1</sup> П	43007.26=	19.15*	7.2281 *	0.0041
(4) <sup>3</sup> Π	50227.78*	49.85*	3.8067*	0.0149
(4) <sup>1</sup> Π	50502.01 *	52.22*	3.8018*	0.0149
(2) <sup>3</sup> Σ <sup>-</sup>	64449.87*	103.98*	2.9190*	0.0253*

By comparing our outcomes of spectroscopic constant  $\omega_e$  with those given in the literature, we obtain a good agreement with the relative differences  $\Delta \omega_e/\omega_e = 7.13\%$ ,  $\Delta R_e/R_e = 9.7\%$ , and  $\Delta B_e/B_e = 0.08\%$  for the ground electronic state  $X^1\Sigma^+$  of TII molecule, [4]. These relative differences

become larger for  $\Delta \omega_e/\omega_e = 12.7$  %,  $\Delta R_e/R_e = 12.0$ %, and in very good agreement with  $\Delta B_e/B_e = 0.29$ % by comparing our values with those of the excited electronic state (1)<sup>3</sup> $\Sigma^+$ , [4].

Similarly, for (1)<sup>3</sup> $\Delta$  electronic state, a strong agreement is obtained between our obtained results and published data for the given spectroscopic constants  $\Delta \omega_e/\omega_e = 5.12\%$ ,  $\Delta R_e/R_e = 1.12\%$ , and  $\Delta B_e/B_e = 4.50\%$ . The spectroscopic constants for the other electronic states are not calculated here since they are unbound states, [4].

#### 3.2 Potential Energy Curves

Based on the 21 valence electrons of the Tl atom using the ECP 60 MDF basis set, and the seven valence electrons of the Iodine atom using the ECP46MDF basis set, 16 singlet and triplet electronic states are investigated and plotted in Figure 1 and Figure 2 as a function of internuclear distance R in the ranges  $1.8\text{\AA} \le \text{R} \le 6.80\text{\AA}$  and  $2\text{\AA} \le$  $\text{R} \le 6\text{\AA}$ , respectively. The potential energy curves, including spin-orbital coupling, are presented in Figure 3 and Figure 4.



Fig. 1: The potential energy curves of the singlet electronic states of the TII molecule without spinorbit coupling



Fig. 2: The potential energy curves of the triplet electronic states of the TII molecule without spin-orbit coupling



Fig. 3: The potential energy curves of the singlet electronic states of the TII molecule with spin-orbital



Fig. 4: The potential energy curves of the triplet electronic states of the TII molecule with spin-orbital

Volume 14, 2023

The ground state of the considered molecule TII has a deep well, while the other investigated electronic states are shallow or unbound. Mainly most of the studied electronic states reach their dissociation limit at around 5Å. The potential energy curve's deepness indicates the molecule's stability, while small forces bind the two atoms for a shallow potential energy curve.

To prove the credibility of our investigated values of the TII molecule, we show in Table 3 the trend of the spectroscopic parameters between the TIBr molecule and our obtained values for the TII molecule, [4]. From bromine to iodine in the Periodic Table, the rotational constant  $B_e$  decreases as the reduced mass increases. This Table also confirms the increase in bond length  $R_e$  and the decrease of harmonic frequency  $\omega_e$  with the decrease of the electronegativity for all the considered electronic states, which can confirm the reliability of our work.

Table 3. The trend study of the spectroscopic constants of the different electronic states of the molecules TlBr and TlI

States	constants	TIBe	TH	trend
$\mathbf{X}_1 \Sigma_{\tau}$	T <sub>e</sub> (cm <sup>-1</sup> )	0*	0.0	
	R.(Å)	2.600*	2.920*	1
	$\omega_{*}(\mathrm{cm}^{-1})$	195.00*	136.00 <sup>b</sup>	5
	B.(cm <sup>-1</sup> )	0.0430*	0.0252 %	5
$(1)^{j}\Sigma^{*}$	T. (cm <sup>-1</sup> )	31174.00*	23391.00*	
	R.(Å)	3.483*	4.1451	2
	$\omega_*(cm^{-1})$	53.00*	28.50 <sup>+</sup>	5
	B.(cm <sup>-1</sup> )	0.024*	0.0125*	×
(1) <sup>7</sup> Δ	T. (cm <sup>-1</sup> )	31611.00*	23730.00*	N .
	R.(Å)	3.570*	37.360 <sup>b</sup>	1
	m <sub>a</sub> (cm <sup>-1</sup> )	47.00*	38.90*	
	B.(cm <sup>-1</sup> )	0.0234*	0.0153 <sup>b</sup>	5
(I) <sup>1</sup> Δ	T <sub>e</sub> (cm <sup>-1</sup> )	31664.00*	23763.11*	5
	R.(Å)	3.616*	4.4173 <sup>b</sup>	
	$\omega_*(\text{cm}^{-1})$	45*	17.55 <sup>±</sup>	×
	B <sub>s</sub> (cm <sup>-1</sup> )	0.230*	0.1103 %	5
(1) <sup>1</sup> Σ <sup>-</sup>	T. (cm <sup>-1</sup> )	31882.00 <sup>±</sup>	23888.61*	5
	R.(Å)	3.654*	4.6182 5	1
	$\omega_{*}(\mathrm{cm}^{-1})$	42*	12.63 <sup>b</sup>	5
	B <sub>e</sub> (cm <sup>-1</sup> )	0.022*	0.0101 5	1
(1)92-	T. (cm <sup>-1</sup> )	31907*	23881.00*	
	R.(Å)	3.666*	4.661*	
	a, (cm <sup>-1</sup> )	42*	11.86°	5
	B <sub>i</sub> (cm <sup>-1</sup> )	0.022*	0.0098%	5
(2) <sup>1</sup> Σ+	$T_{s}(cm^{-1})$	31932*	23886.75*	5
	R.(Å)	3.879*	4.9365*	1
	$\omega_{e}(\text{cm}^{-1})$	334	10.24*	
	B.(cm <sup>-1</sup> )	0.020*	0.0088 h	

<sup>a</sup> [4], <sup>b</sup>Present work

#### **3.3 Permanent Dipole Moment Curves**

By taking the iodine atom at the origin, the curves with and without the spin-orbit of the static dipole moment (DMCs) of the TII molecule are plotted as a function of internuclear distance R in Figure 5, Figure 6, Figure 7 and Figure 8. These DMCs are essential in constructing the molecular orbit bonding model, specifying the polarity and the interaction of electronic states, [17], [18], [19]. One can notice that all the investigated DMCs tend to zero at a large internuclear distance, resulting in a neutral fragment Figure 5 and Figure 8 except for the two electronic states  $(3)^{1}\Sigma^{+}$  and  $(1)^{1}\Delta_{1}$ . These curves tend to have negative values in the negative region within 5.20Å  $\leq R \leq 7.20$  Å (Figure 5 and Figure 7) where the molecule dissociates in the ionic fragment.



Fig. 5: The dipole moment curves as a function of internuclear distance for the singlet electronic states of TII without spin-orbit coupling



Fig. 6: The dipole moment curves as a function of internuclear distance for the triplet electronic states of TII without spin-orbit coupling



Fig. 7: The dipole moment curves as a function of internuclear distance for the singlet spin-orbit coupling electronic states of the TII molecule



Fig. 8: The dipole moment curves as a function of internuclear distance for the triplet spin-orbit coupling electronic states of the TII molecule

## 4 Conclusion

Using an ab initio calculation (CASSCF/MRCI + Q) we investigated in the present work the PECs and the DMCs for 19 and 10 electronic states (singlet and triplet) states of TII molecule. In this calculation, the used basis sets are the ECP60MDF [12] and ECP46MDF [13] for TI and I atoms, respectively. The ground state is confirmed as  $X^{1}\Sigma^{+}$  with deep potential energy curves for the molecule TII. A good agreement can be noticed by comparing our calculated spectroscopic constants in Table 1 with those in the literature. Our previous work on the iodine compounds confirmed the candidacy of the two molecules CaI and BaI for Doppler laser

cooling. This study of laser cooling for the molecule TII has not been done since the first condition, which is that the small difference between the  $R_e$  of the ground and one bound excited state is not verified.

References:

- Metz B, Schweizer M, Stoll H, Dolg M, Liu W. A small-core multiconfiguration Dirac-Hartree-Fock-adjusted pseudopotential for Tl-application to TIX (X = F, Cl, Br, I). *Theor Chem Acc.* 2000; 104: 22-8.
- [2] Goudarzi M, Ghanbari D, Salavati-Niasari M. Photo-catalyst thallium sulfide: characterization synthesis and optical morphologies different of Tl2S nanostructures. Journal of Materials Science: Materials in Electronics. 2015; 26: 8798-806.
- [3] Saoudel A, Amira A, Boudjadja Y, Mahamdioua N, Varilci A, Terzioglu C, Soylu N, Altintas SP. The effect of TIF3 substitution on the weak links behavior and A.C. susceptibility of Bi1.8-xTlxPb0.35Sr2Ca2Cu3OyF3x superconductors. Cryogenics. 2018; 93: 26-33
- [4] Zou W, Liu W. Comprehensive ab initio calculation and simulation on the low-lying electronic states of TlX (X= F, Cl, Br, I, and At). *Journal of computational chemistry*. 2009 Mar;30(4):524-39.
- [5] Zhang X, Zhai H, Liu Y, Sun J. Extensive ab initio calculation on low-lying excited states of CCl+ including spin-orbit interaction. *Journal of Quantitative Spectroscopy and Radiative Transfer.* 2013 Apr 1;1 19: 23-31.
- [6] Li DY, Yang CL, Sun ZP, Wang MS, Ma XG. Theoretical study on the spectroscopic properties of the low-lying electronic states and the laser cooling feasibility of the CaI molecule. *Journal of Quantitative Spectroscopy and Radiative Transfer.* 2021 Aug 1;270:107709.
- [7] Miliordos E, Papakondylis A, Tsekouras AA, Mavridis A. All-electron first principles calculations of the ground and some low-lying excited states of BaI. *The Journal of Physical Chemistry A*. 2007 Oct 4;111(39):10002-9.
- [8] Yang QS, Gao T. The feasibility of laser cooling: an investigation of ab initio of MgBr, MgI and MgAt molecular.

*Spectrochimica Acta Part A:* Molecular and Biomolecular Spectroscopy. 2020 Apr 15;231:118107.

- [9] Madi A., El-Kork N., Zeid I., Korek M., Laser cooling and electronic structure of Be halide anions BeX<sup>-</sup> (X = Cl, Br, F, and I), J. *Chem. Phys.*, 2022; 157:024104.
- [10] Gabedit Allouche AR. a graphical user interface for computational chemistry software. *J Comp. Chem.* 2011; 32:174–82.
- Werner HJ, Knowles PJ, Lindh R, Manby [11] FR, Schütz M, Celani P, Korona T, Rauhut G, Amos RD, Bernhardsson A, Berning A, Cooper DL, Deegan MJO, Dobbyn AJ, Eckert F, Hampel C, Hetzer G, Lloyd AW, McNicholas SJ, Meyer W, Mura ME, Nickla A, Palmieri BP, Pitzer R, Schumann U, Stoll H, Stone AJ. MOLPRO is a package of ab initio programs Molpro: a general-purpose quantum chemistry program package, Werner, 2012, WIREs Computational Molecular Science, Wiley Library, Online https://doi.org/10.1002/wcms.82.
- [12] Y. Wang, M. Dolg, Pseudopotential study of the ground and excited states of Yb2, *Theor. Chem. Acc.* 100 (1998) 124-133
- [13] Lim IS, Schwerdtfeger P, Metz B, Stoll H. All-electron and relativistic pseudopotential studies for the group 1 element polarizabilities from K to element 119. J *Chem Phys* 2005; 122:104103.
- [14] Liu Y, Yuan X, Xiao L, Xu H, Yan B. Electronic structure and spectroscopy of the low-lying electronic states of thallium fluoride: MRCI+ Q study including spinorbit coupling. *Journal of Quantitative Spectroscopy and Radiative Transfer*. 2020 Mar 1; 243:106817.
- [15] Xiao H, Yang QS, Zhu J, Gao T. Ab initio investigation on the low-lying electronic states of thallium bromide. *Spectrochimica Acta Part A*: Molecular and Biomolecular Spectroscopy. 2021 Feb 5; 246:118998.
- [16] Patchkovskii S. Ab initio investigation of potential energy curves of the 23 electronic states of IBr correlating to neutral 2 P atoms. *Physical Chemistry Chemical Physics*. 2006;8(8):926-40.
- [17] Wan MJ, Jin CG, Yu Y, Huang DH, Shao JX. Potential energy curves, transition dipole moments, and radiative lifetimes of KBe molecule. *Chinese Physics B*. 2017; 26:033101.

- [18] Teule JM, Mes J, Bulthuis J, Janssen MH, Stolte S. Laser-Induced Fluorescence Studies of Excited Sr Reactions. 1. Sr (3P1) + H.F. *The Journal of Physical Chemistry* A. 1998; 102: 94828.
- [19] Stradella OG, Castro EA, Fernandez FM. Molecular dipole moment charges from a new electronegativity scale. *Inorganic Chemistry*. 1985; 24:3631-4.

#### **Contribution of Individual Authors to the Creation of a Scientific Article (Ghostwriting Policy)**

- Ghina Chamieh: Conception, design, calculation, and writing the paper.
- Lokman Awad: Revising the paper critically for important intellectual content.
- Mahmoud Korek: Supervised and approved the final version.

#### **Sources of Funding for Research Presented in a Scientific Article or Scientific Article Itself** No funding for this manuscript.

#### **Conflict of Interest**

The authors declare no competing interests.

#### Data availability

All data generated or analyzed during this study are included in this published article.

# Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

https://creativecommons.org/licenses/by/4.0/deed.en US